

RESEARCH ARTICLE

Relation of some biochemical parameters in Van (Turkey) cats with sex, age, eye colour and hair length

Vahdettin ALTUNOK^{1*}, Nuri BAŞPINAR¹, Pınar PEKER AKALIN², Nazmi YÜKSEK³

¹Department of Biochemistry, Faculty of Veterinary Medicine, Selcuk University, Campus, 42075, Konya, Turkey

²Department of Biochemistry, Faculty of Veterinary Medicine, Mustafa Kemal University, Campus, 31060, Hatay, Turkey

³Department of Internal Medicine, Faculty of Veterinary Medicine, Yuzuncu Yıl University, Van, Turkey

Received: 17.10.2017, Accepted: 25.12.2017

*valtunok@selcuk.edu.tr

Van kedilerinde bazı biyokimyasal parametreler ile cinsiyet, yaş, göz rengi ve tüy uzunluğu arasındaki ilişkiler

Eurasian J Vet Sci, 2018, 34, 1, 19-24

DOI:10.15312/EurasianJVetSci.2018.175

Öz

Amaç: Van kedileri, doğu Türkiye kaynaklıdır ve ulusal kültürel varlıkların bir parçasıdır. Van kedilerinin en belirgin özelliği, gözlerinin renkleridir (diskromatopsi). Bu çalışmanın amacı, Van kedilerinde plazma aspartat aminotransferaz, alanin aminotransferaz aktiviteleri, total protein, albümin, glikoz, trigliserit, LDL kolesterol, HDL kolesterol, kan üre / nitrojen ve kreatinin düzeylerini saptamak ve bu parametrelerin cinsiyet, yaş, göz rengi ve tüy uzunluğu ile ilişkisini araştırmaktır.

Gereç ve Yöntem: Bu çalışmada, standart kedi yemi ile beslenen, sağlıklı, 47 adet Van kedisi (Van Kedi Evi, Yuzuncu Yıl Üniversitesi, Van) kan plazması kullanıldı. Kediler, cinsiyet (dişiler-erkek), yaş (1-2; 3-4; 5'in üzerinde), göz rengi (amber-amber, amber-mavi, mavi-amber, mavi-mavi, soldan sağa açıklanan gözler) ve tüy uzunluğu (kısa; uzun) temel alınarak gruplara ayrıldı. Plazma biyokimyasal parametreleri ticari kitlerle kolorimetrik olarak spektrofotometre ile belirlendi.

Bulgular: Cinsiyete ilişkin olarak parametreler arasında fark yoktu. Kısa tüylü kedilerde uzun tüylü kedilere göre albumin düzeyleri daha yüksekti ($p<0.05$). Göz rengi açısından değerlendirildiğinde, mavi-mavi gözlü kedi plazma glikoz düzeyleri diğer göz rengi gruplarına göre çok daha yüksekti ($p<0.001$). Yaş gruplarında, 1-2 yaşındaki kedilerde kreatinin düzeyleri diğer yaş gruplarına göre daha düşüktü ($p<0.05$).

Öneri: Sonuçlarımız, Van kedilerinin plazma biyokimyasal parametrelerinin bazılarının göz rengi, yaş ve tüy uzunluğu ile ilişkili olabileceğini göstermiştir. Ayrıca Van kedilerinden elde edilen bu biyokimyasal bulguların, klinik tanıda yarar sağlayacağı düşünülmektedir.

Anahtar kelimeler: Van Kedileri, Biyokimyasal Parametreler, Göz Rengi, Tüy Uzunluğu

Abstract

Aim: Van cats originate from eastern Turkey and they are a part of the national cultural assets. The most striking characteristic feature of the Van cats is the colour of their eyes (dischromatopsy). The aim of this study was to determine plasma aspartate aminotransferase, alanine aminotransferase activities, total protein, albumin, glucose, triglyceride, total cholesterol, LDL cholesterol, HDL cholesterol, urea/nitrogen, and creatinin levels and the relation of these parameters with sex, age, eye colour and hair length in Van cats.

Materials and Methods: Healthy blood plasma, 47 Van cats (Van Cat House, Yuzuncu Yıl University, Van), fed with standart cat food, were used. Cats were divided into groups based on sex (female-male), age (1-2; 3-4; above 5), eye colour (amber-amber, amber-blue, blue-amber, blue-blue, eyes described from left to right) and hair length (short; long). Plasma biochemical parameters were determined colorimetrically by spectrophotometry, with commercial kits.

Results: There was no difference between the parameters regarding with sex. Albumin levels were higher in the short-haired cats compared to the long-haired cats ($p<0.05$). As regards eye colours, blue-blue eyed cat plasma glucose levels were much higher from than in the other eye colour groups ($p<0.001$). In age groups, creatinin levels were lower ($p<0.05$) in 1-2 year-old cats compared to the other age groups.

Conclusion: Our results indicate that some of the plasma biochemical parameters of the Van cats may be related to their eye-colours, age and hair length. It also can be considered that the biochemical parameters obtained in the Van cat may be useful for the clinical diagnosis.

Keywords: Van cat, Biochemical parameters, Eye-colour, Hair length

Introduction

Van cat which originate from eastern Turkey is a part of our national cultural assets. The most striking characteristic feature of the Van cats is the colours of their eyes. Some of them have different colour of one eye from the other (dischromatopsy) (Gure 1993). In contrast to the other cat species, the Van cats like swimming and playing in water. However, this species is facing with the danger of extinction. Some preventive measures have been taken in order to protect this species (Anonymous 1). Biochemical parameters will be revealed within the scope of protection of the Van cats and important information will be obtained in terms of diagnosis and prognosis of diseases including physiological state changes (pregnancy etc.). Since, to determine the biochemical values of races are more difficult, species specific reference values are generally used. However, it is considered that the most appropriate approach is to determine the reference intervals in such a way as to represent the race (Arıkan et al 2001, 2003, Nisbet 2006, Altunok et al 2007, Macun et al 2010, Şimşek et al. 2015b, Özkan et al 2016).

Biochemical studies of this race have been limited to a few studies in the literature review (Arıkan et al 2003, Macun et al 2010, Altunok et al 2011). In a study on blood groups, A blood group was found to be 40% and B blood group was found to be 60% in the Van cats (Arıkan et al 2003). In the same study, the percentages of blood groups A and B in the Angora cats were determined as 53.6% and 46.4%, respectively and it was revealed that there are differences between races. In another study of genetically differentiating the Van cats from other cat breeds, Altunok et al (2011) reported that phosphogluconate dehydrogenase, malic enzyme and esterase D loci are different between Van cats and other cat breeds. Macun et al (2010) determined the some biochemical parameters of the Angora and Van cats during their gestation and lactation period. The ALT, AST activities, glucose, cholesterol, albumin, Pi and Mg levels were within the reference range but total protein, globulin and Ca levels were above the reference range according to other cat breeds. When examined in terms of biochemical parameters such as serum AST, ALT, lactate dehydrogenase (LDH), creatine kinase (CK) activities, glucose, urea, creatinine, potassium (K), chlorine (Cl) and sodium (Na) levels in female, male, and under 1 year old kittens were determined within the same reference range as other cat breeds (Sönmez and Ağaoğlu 2010). Akkan et al (2005) reported that serum activities of AST, ALT, ALP, and levels of bilirubin in the Van cats were the same reference values of other cat breeds. Yörük (2000) determined the total cholesterol, triglyceride and total protein concentrations as mg/dl by the colorimetric method and also the levels of VLDL, LDL and HDL as percent (%) by electrophoretic method in 10 Van cats. Altunok et al (2007) determined that some trace element levels in the male cats were different from than in the female cats in this study of the Van cats and also levels of aluminum, copper, manganese, strontium and zinc were higher in blue-blue eyed cats than in the other different coloured eyed cats ($p < 0.05$).

Although some biochemical parameters were revealed (Macun et al 2010, Sönmez and Ağaoğlu 2010, Karaca 2009, Söyler 2006, Özkan et al 2016) in the Van cats, it was observed that these parameters were not related to age, gender, eye colour and hair length in the literature reviews except Altunok et al (2007) and Özkan et al (2016). It is reported that blood biochemical parameters may change under different factors such as sex, age, race, climate, stress, and pregnancy (Eksen et al 1992, Awah and Nottidge 1998, Strasser et al 2001, Altunok et al 2007, Uslu et al 2012, Şimşek et al 2015a,b). In different age groups of cattle, plasma ALT, AST and alkaline phosphate (ALP) activities were found different (Sharma and Bisoi 1995). In Angora cats, and plasma creatinine levels reported to increased with age. In Kangal dogs, cholesterol levels were lower and total protein levels were higher in adults compared to puppies (Çınar et al 2010). Also, Arıkan et al (2001) reported that biochemical parameters differ within different race of sheep.

Therefore, the aim of this study was to investigate the relationship of plasma aspartate aminotransferase, alanine aminotransferase activities, total protein, albumin, glucose, triglyceride, total cholesterol, LDL cholesterol, HDL cholesterol, blood urea/nitrogen, and creatinin levels with age, gender, eye colour and hair length in the Van cats.

Materials and Methods

To determine selected biochemical parameters of the Van cats, 47 healthy animals were used (Van cat house, YuzuncuYıl University, Van, Turkey). The cats were fed with standard canned food (LA-CAT, Israel). Cats were divided into four groups based on age (1 - 2; 3 - 4; and > 5 years old), sex (female; male), eye colour (amber-amber; amber-blue; blue-amber and blue-blue, eyes described from left to right) and hair length (short; long). Blood samples were collected from each cats in the morning (10:00 a.m.) by puncture of the cephalic vein.

Plasma (EDTA) samples were separated by centrifugation (3000 rpm, 15 minutes), and stored at -80 °C until analysis. Plasma aspartate aminotransferase (AST), alanine aminotransferase (ALT) activities, total protein, albumin, glucose, triglyceride, total cholesterol, LDL cholesterol, HDL cholesterol, blood urea/nitrogen, and creatinin levels were determined colorimetrically (UV 2100 UV-VIS Recording Spectrophotometer Shimadzu, Japan) with commercial kits (Human Diagnostics Worldwide, GmbH, Germany).

Statistical analyses

All the values are expressed as mean \pm SE. The results of groups were analyzed by Student's t-test (sex and hair length groups) and ANOVA (age and eye colour) with posthoc Duncan multiple range test (SPSS for Windows, release 12.0). In all cases, probability of error less than 0.05 was chosen as the criterion of

statistical significance.

Results

Biochemical parameters and the relations of these parameters with the variables (sex, age, eye colour and hair length) in the Van cats are given in Table 1, 2, 3 and 4.

As regards sex, there was not any statistically significant difference in the parameters (Table 1). When assessed for age, creatinin levels were lower ($p<0,05$) in 1-2 year-old group compared to 3-4 year-old and above 5 year-old cats (Table 2).

Blue-blue eyed cats had much higher ($p<0,001$) plasma glucose levels than in the other eye colour groups (Table 3). Plasma albumin levels were higher ($p<0,05$) in the short haired cats compared to the long haired cats (Table 4). There was not any statistically significant difference between the other biochemical parameters and the variables.

Discussion

In the present study, some biochemical parameters of the Van cats in relation to their sex, age, eye colour and hair length have been shown in the tables 1, 2, 3, and 4.

Table 1. Biochemical parameters of the Van cats according to the sex differences (Mean \pm SE)

Sex	Female n=31	Male n=16	Total n=47	References
AST (U/L)	21.55 \pm 2.13	20.20 \pm 2.17	21.10 \pm 1.58	26-43 (Kaneko 1997), 10-100 (Anonymous 2)
ALT (U/L)	20.66 \pm 1.54	19.42 \pm 1.51	20.25 \pm 1.14	6-83 (Kaneko 1997), 10-100 (Anonymous 2)
Total Protein (g/dl)	7.23 \pm 0.15	6.70 \pm 0.24	7.05 \pm 0.013	6.3-10.3g/dl (Lehmann et al. 1997, Eliotta and Barber 1998, Reynolds et al. 2010)
Albumin (mg/dl)	2.93 \pm 0.07	2.78 \pm 0.09	2.88 \pm 0.05	2.6-3.9 mg/dl (Reynolds et al. 2010)
Glucose (mg/dl)	116.92 \pm 10.81	94.45 \pm 6.99	109.43 \pm 7.68	73-134 (Kaneko 1997), 61-170 (Anonymous 2, Reynolds et al. 2010)
Triglyceride (mg/dl)	49.22 \pm 6.25	50.11 \pm 8.39	49.52 \pm 4.96	35-62 mg/dl (Lehmann et al. 1997)
Total Cholesterol (mg/dl)	71.19 \pm 6.83	68.48 \pm 5.90	70.29 \pm 4.92	69.50-180 mg/dl (Eliotta and Barber 1998)
LDL (mg/dl)	40.67 \pm 3.16	39.88 \pm 4.23	40.41 \pm 2.51	11-59 (Demacker 1997, Watson 1995)
HDL (mg/dl)	19.56 \pm 2.37	25.70 \pm 3.63	21.61 \pm 2.01	71-148 (Demacker 1997, Watson 1995)
				37.40 (Mosallanejad et al. 2016)
BUN (mg/dl)	20.91 \pm 0.95	19.81 \pm 1.05	20.55 \pm 0.72	20-30 (Kaneko 1997)
Creatinin (mg/dl)	1.25 \pm 0.05	1.032 \pm 0.09	1.28 \pm 0.05	0.9-1.9 (Kaneko 1997)

There was not any statistical significant differences ($p>0.05$)

Table 2. Biochemical parameters of the Van cats according to the age of years differences (Mean \pm SE)

Age	1-2 years n=21	3-4 years n=17	>5 years old n=9
AST (U/L)	21.45 \pm 2.27	23.65 \pm 3.19	16.04 \pm 2.17
ALT (U/L)	19.40 \pm 1.64	20.28 \pm 2.31	22.16 \pm 1.94
Total Protein (mg/dl)	7.10 \pm 0.22	6.94 \pm 0.21	7.13 \pm 0.24
Albumin (mg/dl)	2.92 \pm 0.09	2.80 \pm 0.09	2.91 \pm 0.09
Glucose (mg/dl)	120.90 \pm 14.53	104.74 \pm 7.33	90.50 \pm 12.14
Triglyceride (mg/dl)	59.26 \pm 9.70	42.23 \pm 4.74	38.94 \pm 3.18
Total Cholesterol (mg/dl)	72.34 \pm 9.76	64.30 \pm 4.33	75.47 \pm 6.68
LDL (mg/dl)	37.96 \pm 3.96	38.24 \pm 4.08	49.74 \pm 4.34
HDL (mg/dl)	23.81 \pm 3.76	18.25 \pm 2.37	22.09 \pm 3.02
BUN (mg/dl)	20.95 \pm 1.20	19.04 \pm 1.10	21.82 \pm 1.29
Creatinin (mg/dl)	1.15 \pm 0.06 a	1.39 \pm 0.07 b	1.40 \pm 0.12 b

Differences in the same row are statistically significant ($p<0.05$)

Table 3. Biochemical parameters of the Van cats according to the eye colour differences (Mean \pm SE) Eye colours (left eye - right eye)

Eye Colour	Amber-Amber n=13	Amber-Blue n=12	Blue-Amber n=12	Blue-Blue n=10
AST (U/L)	18.45 \pm 1.86	26.13 \pm 4.51	17.77 \pm 2.54	22.56 \pm 3.52
ALT (U/L)	18.96 \pm 1.90	21.29 \pm 2.98	20.94 \pm 1.78	18.93 \pm 2.85
Total Protein (mg/dl)	7.10 \pm 0.19	6.68 \pm 0.22	7.50 \pm 0.26	7.02 \pm 0.34
Albumin (mg/dl)	2.81 \pm 0.04	2.79 \pm 0.13	2.93 \pm 0.08	2.95 \pm 0.17
Glucose (mg/dl)	114.47 \pm 10.52 ^a	84.66 \pm 4.74 ^a	83.99 \pm 3.83 ^a	162.87 \pm 27.91 ^b
Triglyceride (mg/dl)	45.87 \pm 2.58	50.93 \pm 10.50	39.53 \pm 7.51	67.05 \pm 18.28
Total Cholesterol (mg/dl)	77.19 \pm 14.63	70.06 \pm 7.46	67.91 \pm 6.70	66.42 \pm 9.64
LDL (mg/dl)	39.75 \pm 3.30	39.71 \pm 5.80	45.88 \pm 5.23	36.29 \pm 6.46
HDL (mg/dl)	19.57 \pm 2.15	20.00 \pm 3.28	25.35 \pm 3.80	22.37 \pm 7.44
BUN (mg/dl)	21.75 \pm 1.33	19.50 \pm 0.67	18.45 \pm 1.76	22.69 \pm 1.98
Creatinin (mg/dl)	1.39 \pm 0.09	1.20 \pm 0.08	1.26 \pm 0.12	1.24 \pm 0.09

Differences in the same row are statistically significant (p<0.001)

Table 4. Biochemical parameters of the Van cats according to the hair length differences (Mean \pm SE)

Hair-length	Short-haired n=32	Long-haired n=15
AST (U/L)	20.71 \pm 1.97	21.96 \pm 2.71
ALT (U/L)	19.99 \pm 1.31	20.81 \pm 2.30
Total Protein (mg/dl)	7.10 \pm 0.18	6.95 \pm 0.13
Albumin (mg/dl)	2.95 \pm 0.07 ^a	2.71 \pm 0.05 ^b
Glucose (mg/dl)	112.90 \pm 9.87	101.74 \pm 11.75
Triglyceride (mg/dl)	53.90 \pm 6.97	39.81 \pm 3.00
Total Cholesterol (mg/dl)	72.99 \pm 6.83	64.31 \pm 4.65
LDL (mg/dl)	40.87 \pm 3.11	39.38 \pm 4.33
HDL (mg/dl)	23.02 \pm 2.73	18.49 \pm 2.23
BUN (mg/dl)	20.55 \pm 0.96	20.58 \pm 0.94
Creatinin (mg/dl)	1.27 \pm 0.06	1.32 \pm 0.06

Differences in the same row are statistically significant (p<0.05)

Determination of AST and ALT activity informs about liver and soft tissue damage. In this study plasma activities of AST and ALT were consistent with the levels of those reported by Sönmez and Ağaoğlu (2010) and Macun et al (2010), whereas Özkan et al (2016) reported higher levels of ALT levels in Van cats. There was not any association between the variables (sex, age) and AST, ALT activities, which was in good agreement with Özkan et al (2016) who reported unchanged AST and ALT activities between sex and age in Van cats.

As regards total protein and albumin levels, it was found that levels were slightly lower from than in Macun et al (2010) who reported in the Van cats, whereas the results of this study were

in good agreement with Özkan et al (2016) and cat reference values reported by Reynolds et al (2010). There was not any correlation between total protein and variables in this study, but it was observed that albumin levels in the short-haired group were higher than in the long-haired group (p<0.05). There was not any literature on the relationship between hair length and albumin levels. It is suggested to investigate this relation detailed.

In the literature, there is discrepancy in plasma glucose levels, such as; plasma glucose levels were 109,43 \pm 7,68 mg/dl in our study, Sönmez and Ağaoğlu (2010) reported plasma glucose levels as 86-87 mg/dl and Özkan et al reported as 58.0 \pm 3.2 mg/dl.



Nevertheless, our results were consistent with the values reported in cats (Kaneko 1997, Anonymous 2, Reynolds et al 2010). In the present study, it was determined that plasma glucose levels were not related to the variables except eye colour, that is; glucose levels were found higher in the Van cats with blue-blue eyed than in the other eye colour groups ($p<0.001$). There was not information about the relationship between eye colour and plasma glucose levels. Similar to this novel result, Altunok et al (2007) determined that the concentrations of serum Al, Cu, Mn, Sr and Zn in the blue-blue eyed cats were higher than the cats with amber-amber, amber-blue and blue-amber coloured eye ($p<0.05$). Further studies may be established to elaborate the carbohydrate metabolism in the Van cats and to elucidate the link between carbohydrate metabolism and genes that determine eye colour.

It is important to reveal plasma lipid profile in terms of the investigation of obesity and follow-up of metabolic syndrome in cats. In the present study, plasma triglyceride levels were compatible with Yörük (2000), whereas total cholesterol levels were slightly lower. Nevertheless, total cholesterol levels were consistent with the reference values reported in cats (Reynolds et al 2010).

Although, Yörük (2000) represented blood VLDL, LDL ve HDL levels by electrophoretic method and reported to 10.15%, 20.05%, 69.67%, respectively, there was not any study reporting plasma HDL and LDL levels as mg/dl in the Van cats. For this reason, we believe that, it is important to establish blood lipid profile of the Van cats in terms of follow-up of related diseases. While LDL levels were compatible with the values reported for the other cat breeds (Kaneko 1997), HDL levels were lower than the other cat breeds (Demacker 1997, Watson 1995, Mossallanejad et al 2016). The low HDL levels in the Van cats may be a specific level for the race of the Van cats. It is suggested that the subject of the low HDL levels in the Van cats should be investigated with more detailed studies.

Blood urea/nitrogen and creatinine levels are two important parameters used to follow-up of kidney diseases. In this present study, the blood urea/nitrogen values obtained in the Van cats were consistent with the values stated by Özkan et al (2016), whereas levels were slightly lower than the levels reported by Sönmez and Ağaoğlu (2010) in the Van cats. The blood urea/nitrogen values and the variables in this study were not correlated which the result was in good agreement with the above researchers (Sonmez and Ağaoğlu 2010, Özkan et al 2016). Creatinin levels were consistent with values reported in the Van Cats (Macun et al 2010, Özkan et al 2016) and the other cat breeds (Kaneko 1997). Özkan et al (2016) reported higher ($p<0.001$) creatinin levels in male Van cats compared to females, furthermore, 12-24 month and 24 to more-month male cats had higher creatinin levels compared to female cats at the same ages. In our study, creatinin levels were not different between the sexes, whereas the levels were higher in 3 year-old cats than in 1-2

year-old cats ($p<0.05$). The study (Şimşek et al 2015b), established for the determination of age relationship with urea and creatinine levels in the Angora cats, showed that blood urea levels did not change with age and plasma creatinine levels which increased with age. In our study, we also could not find any relationship with BUN and the variables, whereas creatinine levels of over 3 year-old cats were higher than in 1-2 year-old cats ($p<0.05$). It was obtained that, the results of parameters (blood urea/nitrogen and creatinin levels) in our study were consistent with the study in the Angora cats (Şimşek et al 2015a).

Conclusion

Some biochemical parameters (plasma HDL, LDL levels) in the Van cats were presented for the first time. Plasma glucose levels in the blue-blue eyed cats were found higher than in the other eyed cats. Albumin levels of the short-haired cats were higher than in the long-haired cats and creatinin levels were higher in the old cats than young cats. It can be suggested to investigate these relations in the Van cats and verify with more observations based on a greater number of cats. It also can be considered that the biochemical parameters obtained in the Van cat may be useful for the clinical diagnosis.

Acknowledgements

This study was approved by the Selcuk University Local Ethic Committee (29.03.2016/2016-34) and supported by SUBAP (Project Number: 16401063). We appreciate to the personnel of the Van Cat House, Van, Turkey. The abstract of this study was published in the '1st International Veterinary Biochemistry and Clinical Biochemistry, 12-14 April 2018/Hatay/TURKEY' congress book.

References

- Akkan HA, Karaca M, Tütüncü M, 2005. Haemobartonellosis in Van Cats. *Turk J Vet Anim Sci*, 29, 709-712.
- Altunok V, Yazar E, Yüksek N, 2007. Selected blood serum elements in Van (Turkey) cats. *Acta Vet Brno*, 76, 171-177.
- Altunok V, Yüksek N, Berkman CC, Ağaoğlu ZT, Togan İ, 2011. Genetic structure and variation of Van cats. *Biochem Genet*, 49, 511-522.
- Anonymous1. <https://vankedisi.yyu.edu.tr/AkademikBirimler/index.php?s=vankedisi>, Access:13.10.2017
- Anonymous 2./2011 <http://www.2ndchance.info/normal-dogandcatbloodvalues.htm>. Access: 06.08.2017.
- Arıkan Ş, Duru SY, Gurkan M, Agaoglu ZT, Giger U, 2003. Blood type A and B frequencies in Turkish Van and Angora cats in Turkey. *J Vet Med Series A*, 50, 303-306.
- Arıkan Ş, Yiğit A, Akçapınar H, Zengin N, Taşdemir U, 2001. Sakız Karayaka ve Sakız x Karayaka G1 melezi koyunlarının bazı hematolojik değerlerinin karşılaştırılması. *Ankara Üni Vet Fak Derg*, 49: 101-106.



- Awah JN and Nottidge HO, 1998. Serum biochemical parameters in clinically healthy dogs in Ibadan. *Trop Vet*, 16, 123-129.
- Çınar M, Erat S, Arıkan Ş, Mamak N, Oğrak YZ, Güzel M, 2010. Kangal Köpeklerinde Bazı Biyokimyasal Parametreler Üzerine Yaş ve Cinsiyetin Etkisi. *Erciyes Üni Vet Fak Derg*, 7, 109-116.
- Demacker PNM, van Heijst PJ, Hak-Lemmers HLM, 1997. A study of the lipid transport system in the cat. *Felix domesticus. Atherosclerosis*, 66, 113-123.
- Eksen M, Agaoglu ZT, Keskin E, 1992. Some haematological values of healthy cats in Van. *Eurasian J Vet Sci*, 8, 45-47.
- Gure A, 1993. Van Kedisi (Van cat). In, Abdulkadiroglu A, Yigit M, Oguzbasaran B (Ed): *Van Kütüğü*. No: 8, Yüzüncü Yıl University, Van, Turkey, pp; 722-726,
- Kaneko, JJ. Harvey JW, Bruss ML, 1997. *Clinical Biochemistry in Domestic Animals*, 5th Edition, Academic Press, California, USA.
- Karaca S, 2009. Kan kedilerinde gebelik ve laktasyon dönemlerinde iz element ve mineral düzeylerinin araştırılması, Yüksek Lisans Tezi, Sağlık Bilimleri Enstitüsü, Yüzüncü Yıl Üniversitesi, Van.
- Macun H, Çınar M, Erat S, Arıkan Ş, 2010. Ankara ve Van Kedilerinin Gebelik ve Laktasyon Dönemlerine Ait Bazı Biyokimyasal Parametrelerinin Karşılaştırılması. *Erciyes Üniv Vet Fak Derg*, 7, 99-108.
- Mosallanejad B, Avizeh R, Razi Jalali M. Pourmahdi, M, 2016. Comparative evaluation between chitosan and atorvastatin on serum lipid profile changes in hyperlipidemic cats. *Iran J Vet Res*, 17, 36-40.
- Nisbet C, Yarım GF, Çiftçi G, 2006. Sağlıklı Karayaka ırkı konyunlara ait bazı serum biyokimyasal değerleri. *Ankara Üniv Vet Fak Derg*, 53: 57-59.
- Özkan C, Kozat S, Kaya A, Aygöl Y, 2016. Some hematological and biochemical parameter levels in healthy Van cats at different age and gender. *Eurasian J Vet Sci*, 32, 214-219.
- Reynolds BS, Concordet D, Germain CA, Daste T, Boude KG and Lefebvre HP, 2010. Breed Dependency of Reference Intervals for Plasma Biochemical Values in Cats. *J Vet Int Med*, 24, 809-818.
- Sharma M and Bisoi PC, 1995. Clinically important serum enzymes of indigenous cattle. *Ind Vet J*, 72, 21-24.
- Şimşek Ö, Arıkan Ş, Çınar M, 2015a. Reference values for selected hematological and biochemical blood parameters from prepregnancy to advanced gestation in Angora cats. *Turk J Vet Anim Sci*, 39, 29-33.
- Şimşek Ö, Çınar M, Arıkan Ş, 2015b. Changes in selected hematology and serum biochemistry in Turkish Angora cats (*Felis catus*) during growth period. *J Adv Vet Anim Res*, 2(1), 34-39.
- Sonmez N and Ağaoğlu ZT, 2010. Van Kedilerinde Kardiyak Troponin Seviyelerinin Araştırılması. *Van Vet J*, 21, 21-25.
- Soyler M, 2006. Van Kedi eritrositlerinden karbonik anhidraz enziminin (E.C.4.2.1.1.) saflaştırılması ve karakterizasyonu. Yüksek Lisans Tezi, Fen Bilimleri Enstitüsü, Yüzüncü Yıl Üniversitesi, Van.
- Strasser A, Seiser M, Heizmann V, Niedermuller H, 2001. The influence of season on hematological and clinical parameters in a beagle dog colony. *Kleintierpraxis*, 46, 793-804.
- Uslu BA, Şendağ S, Uçar Ö, Gülyüz F, Taşal İ, Çetin N, Wehrend A, 2012. The Effects of Pregnancy, Early Lactation Weeks and the Number of Suckling Kittens on the Serum Mineral and Metabolite Profiles of Van Cats in Breeding Season, *Kafkas Univ Vet Fak Derg*, 18, 691-700.
- Watson TDG, Butterwick RF, McConnell M, 1995. Development of methods for analysing plasma lipoprotein concentrations and associated enzyme activities in the cat and their use to measure the effect of pregnancy and lactation. *Am J Vet Res*, 56, 289-296.
- Yoruk İH, 2000. Evcil hayvanlarda serum lipid, total protein ve lipoprotein düzeylerinin araştırılması, Yüksek Lisans Tezi, Sağlık Bilimleri Enstitüsü, Yüzüncü Yıl Üniversitesi, Van.