



## RESEARCH ARTICLE

### Evaluation of acute phase proteins and cytokines in dogs with parvoviral enteritis

Mahmut Ok<sup>1\*</sup>, Cenk Er<sup>1</sup>, Ramazan Yıldız<sup>2</sup>

<sup>1</sup>Department of Internal Medicine, Faculty of Veterinary Medicine, University of Selcuk, Konya, <sup>2</sup>Department of Internal Medicine, Faculty of Veterinary Medicine, University of Mehmet Akif Ersoy, Burdur, Turkey

Received: 05.03.2015, Accepted: 13.04.2015

\*mok@selcuk.edu.tr

### Parvoviral enteritli köpeklerde akut faz proteinler ve sitokinlerin değerlendirilmesi

Eurasian J Vet Sci, 2015, 31, 3, 143-147  
DOI:10.15312/EurasianJVetSci.2015310970

#### Öz

**Amaç:** Bu çalışmanın amacı parvoviral enteritli köpeklerde akut faz proteinler ve sitokinlerdeki değişimleri değerlendirmektir.

**Gereç ve Yöntem:** Çalışmanın materyalini 2 - 8 aylık 46 parvoviral enteritli (PVE grubu) ve 8 sağlıklı (Kontrol grubu) köpek oluşturdu. Parvoviral enteritli köpeklerde anoreksi, ateş, depresyon, uyuşukluk, kusma ve kanlı ishal belirlendi. Köpeklerde parvoviral enteritis teşhisi dışkı parvovirüs antijen testi ile doğrulandı. Parvoviral enteritis teşhisi konulan köpeklerden kan örnekleri alınarak serum interleükin 1β (IL-1β), tümör nekrozis faktör alfa (TNF-α), gama interferon (INF-γ), C-reaktif protein (CRP), serum amiloid A (SAA), fibrinojen ve protein C (PC) düzeyleri ELISA metoduyla ölçüldü.

**Bulgular:** Sağlıklı grupla karşılaştırıldığında, parvoviral enteritli köpeklerin serum IL-1β, TNF-α, INF-γ, CRP, SAA, fibrinojen ve PC seviyelerinde istatistiksel olarak önemli artışlar belirlendi.

**Öneri:** Parvoviral enteritli köpeklerde serum IL-1β, TNF-α, INF-γ, CRP, SAA, fibrinojen ve PC seviyelerinde artışların belirlenmesi yangının değerlendirilmesinde faydalı olacağı ifade edilebilir.

**Anahtar kelimeler:** Köpek, parvoviral enteritis, akut faz proteinler, sitokinler

#### Abstract

**Aim:** The aim of the present study was to evaluate the alterations in acute phase proteins and cytokines in dogs with parvoviral enteritis.

**Materials and Methods:** Aged between 2 - 8 months, forty-six dogs with parvoviral enteritis (PVE group) and 8 healthy dogs (Control group) were as material in this study. Anorexia, fever, depression, lethargy, vomiting and hemorrhagic diarrhea were determined in the dogs with parvoviral enteritis. Parvovirus infection in dogs was confirmed by examining faeces via parvovirus antigen test. Blood samples were collected from all dogs. Serum interleukin-1β (IL-1β), tumor necrosis factor alpha (TNF-α), interferon-γ (INF-γ), C-reactive protein (CRP), serum amyloid A (SAA), fibrinogen and protein C (PC) concentrations were measured by ELISA.

**Results:** Serum IL-1β, TNF-α, INF-γ, CRP, SAA, fibrinogen and PC concentrations were statistically significantly increased in the dogs with parvoviral enteritis compared to control group.

**Conclusion:** It may be stated that increased serum IL-1β, TNF-α, INF-γ, CRP, SAA, fibrinogen and PC concentrations may be useful in assessing the inflammation in canine parvovirus infection.

**Keywords:** Dog, parvoviral enteritis, acute phase protein, cytokines



## Introduction

Canine parvovirus (CPV) infection is considerable cause of high morbidity and mortality in dogs whelps younger than 6 months (Otto et al 1997, McCaw and Hoskins 2006, Er ve Ok 2015). Canine parvovirus infection has two clinical forms as acute hemorrhagic enteritis and myocarditis. Myocarditis form can develop from infection in utero or in puppies less than 8 weeks old. However acute enteritis form is mostly seen in puppies up to 6 months of age. In myocarditis form, the infected puppies may be found as dead without any symptoms within 24 hours (Prittie 2004, Goddard and Leisewitz 2010). In canine parvoviral enteritis, mortality without treatment has been reported as 91%, however, it is associated with a survival rate as low as 64% with treatment (Kariuki et al 1990).

Serum concentrations of acute phase proteins (APPs) such as C-reactive protein (CRP), serum amyloid A (SAA) and haptoglobin increase in within a few hours following infection (Eckersallet al 2010). The acute phase response assures reliable markers for diagnosing and prognosticating diseases in both people and animals (Kjelgaard-Hansen and Jacobsen 2011). APPs are synthesized in the liver in response to release of proinflammatory cytokines in diseases such as bacterial and viral infections, immuno-mediated disease, neoplasia, tissue injury (trauma), necrosis and burns (Turgut 2000, Murata et al 2004, Eckersallet al 2010, Kocaturket al 2010). Serum APPs levels have been detected to be eight times more sensitive to inflammatory diseases than white blood cells (Ceronet al 2005).

Clinical applications for APPs have been widely demonstrated prognostication as well as for detection of clinical disease and chronic inflammation (Cary 2011). CRP and SAA have been used in the diagnoses of infection in dogs. Furthermore, increased CRP and SAA concentrations have been detected in dogs with systemic inflammation (Gebhardt et al 2009, Jain et al 2011, Langhorn et al 2014). Serum APPs concentrations have significantly increased in dogs with parvoviral enteritis, especially, CRP seems to be a potent predictor of mortality of diseases (Kocaturk et al 2010).

Endotoxemia have been reported in dogs with parvoviral enteritis (PVE). Bloody diarrhea in parvoviral enteritis results from endotoxemia and cytokine production (Otto et al 1997). Bacterial translocation and coliform septicemia are major factors in the mortality associated with PVE (Isogai et al 1989). Acute phase reaction is alerted by the release of cytokines such as IL-1 $\beta$ , IL-6 and TNF- $\alpha$  at the site of inflammatory lesions or infections (Bochsler and Slausan 2002, Carry 2013). IL-1 $\beta$ , IL-6, TNF- $\alpha$  and IFN- $\gamma$  are produced by inflammatory cells. These proinflammatory cytokines induce local and systemic reactions (Jain et al 2011). It is known that sepsis occurs in dogs with PVE. Serum IL-1 $\beta$ , IL-6 and

TNF- $\alpha$  levels are found to be increased in sepsis. IL-6 may be a reliable indicator of the severity of a systemic bacterial infection (Rau et al 2007). It has been noted that serum IL-6 and TNF- $\alpha$  levels might be valuable for evaluating septic patients (Song et al 2012).

The aim of this study was to evaluate the alteration of acute phase proteins and cytokines in dogs with PVE.

## Materials and Methods

The institutional ethical committee approved this prospective study. In this study, 46 dogs with PVE (PVE group) and 8 healthy dogs (Control group) were used. Dogs, aged between 2 and 8 months, were brought into the Faculty of Veterinary Medicine, Internal Medicine Department. Clinical examinations were performed for all dogs. Faeces samples were collected from clinically PVE suspected dogs (anorexia, depression, lethargy, vomiting and hemorrhagic diarrhea) by using rectal swab. Faeces samples were evaluated by the quick parvovirus antigen detection test (CPV Ag test, rapidly test kits, Vettek Medical Istanbul, Turkey).

After the diagnosis of PVE, blood samples were collected from dogs. Blood samples were obtained after centrifugation and stored at -80°C deep freeze until analysis. Canine CRP (Eastbiopharm, China), canine SAA (Eastbiopharm, China), canine protein C (TSZ ELISA, USA) and canine fibrinogen (Eastbiopharm, China) concentrations were measured by ELISA method in Synergy HT multi-mode microplate reader (BioTek Inc, USA) device described by the manufacturer. Measurable sensitivity of CRP is 0.051 mg/L, and test interval of CRP level is 0.1 mg/L and 30 mg/L, measurable sensitivity of SAA is 0.047  $\mu$ g/mL and test interval of SAA level is 0.1  $\mu$ g/mL and 40  $\mu$ g/mL, measurable sensitivity of protein C (PC) is less than 0.15 ng/mL and test interval of PC level is 0.15 ng/mL and 40 ng/mL, and measurable sensitivity of fibrinogen is 0.023 mg/mL and test interval of fibrinogen level is 0.05 mg/mL and 15 mg/mL. Canine IL-1 $\beta$  (Eastbiopharm, China), canine TNF- $\alpha$  (Eastbiopharm, China) and canine IFN- $\gamma$  (Eastbiopharm, China) levels were measured by ELISA method in Synergy HT multi-mode microplate reader (BioTek Inc, USA) device described by the manufacturer.

Measurable sensitivity of IL-1 $\beta$  is 0.1 pg/mL, and test interval of IL-1 $\beta$  level is 0.2 pg/mL and 60 pg/mL, measurable sensitivity of TNF- $\alpha$  is 0.01 ng/L and test interval of TNF- $\alpha$  level is 0.03 ng/L and 9 ng/L and measurable sensitivity of INF- $\gamma$  is 2.35 ng/mL and test interval of INF- $\gamma$  level is 5 ng/L and 1000 ng/L.

All data were presented as mean $\pm$ SEM. Two sample T test was used to determine the differences between groups (SPSS 19.0 for Windows). The statistical significance level was considered to be P<0.05.





Table 1. Concentrations of acute phase proteins in dogs.

Parameters	Control (n:8)	PVE (n: 46)	P value
CRP (mg/L)	2.27±0.19	3.47±0.58	P<0.05
SAA (µg/mL)	2.30±0.52	3.93±0.87	P<0.05
Fibrinogen (mg/mL)	3.08±0.42	4.68±0.65	P<0.05
Protein C (ng/mL)	0.68±0.10	2.10±0.23	P<0.001
IL-1β (pg/mL)	9.30±0.53	12.9±1.24	P<0.009
TNF-α (ng/L)	0.24±0.76	1.04±0.27	P<0.007
INF-γ (ng/L)	104±0.86	176±23.2	P<0.003

PVE: Parvoviral group, CRP: C reactive protein, SAA: Serum amyloid A, IL-1β: Interleukin 1β, TNF-α: Tumor necrosis factor alpha, INF-γ: Interferon-γ.

## Results

Dogs with PVE had anorexia, lethargy, depression, fever, vomiting and hemorrhagic diarrhea. Serum CRP, SAA, fibrinogen, IL-1β, TNF-α, INF-γ and PC concentrations in dog with PVE and healthy are summarized in Table 1. Serum CRP (P<0.05), SAA (P<0.05), fibrinogen (P<0.05), PC (P<0.001), IL-1β (P<0.009), TNF-α (P<0.007) and INF-γ (P<0.003) levels were significantly increased in dogs PVE when compared to healthy dogs.

## Discussion

Parvoviral infection, one of the most important (lethal) diseases among infectious diseases of dogs, is very difficult to treat. Anorexia, depression, lethargy, fever, tachycardia and tachypnea, vomiting, diarrhea (that can change to mucoid or hemorrhagic) are the common symptoms among dogs with PVE (Pollock and Coyne 1993, Ok et al 2000, Yesilbag et al 2007, Yilmaz and Senturk 2007, Goddard and Leisewitz 2010, Kocaturk et al 2010, Er ve Ok 2015). These symptoms were also diagnosed in this study. The diagnosis of this disease in this study was confirmed by parvovirus antigen test.

This study showed the changes of APPs and cytokines in dogs with PVE (Table 1). APPs are synthesized in the liver in response to release of proinflammatory cytokines in diseases such as bacterial and viral infections, immuno-mediated disease, neoplasia, tissue injury (trauma), necrosis and burns (Turgut 2000, Murata et al 2004). These APPs might be beneficial in determining the disease severity and possibly the response to treatment or prognosis, on a disease-specific basis (Murata et al 2004, Ceronet al 2005, Kjelgaard-Hansen and Jacobsen 2011). CRP and SAA are useful parameters among APPs to indicate inflammation in human and animals (Eckersell and Bell 2010). Increased CRP and SAA concentrations have been detected in dogs with systemic inflammation (Fransson et al 2007, Gebhardt et al 2009, Jain et al 2011, Langhorn et al 2014). CRP and SAA have also been

reported to be significantly increased in dogs with PVE (Yule et al 1997, Kocaturk et al 2010). CRP and SAA concentrations significantly increased in dogs with pyometra (Jitpean et al 2014). In present study, CRP, SAA, and fibrinogen concentrations statistically increased in dogs with PVE compared to healthy dogs (Table 1). The reason for this increase in dogs with PVE could be related to an inflammatory process. So, serum CRP, SAA and fibrinogen levels could be important markers for inflammatory process in dogs with PVE. Similarly, several studies have reported that CRP and SAA can be useful markers for diagnosis and prognosis in various diseases (Mastrorilli et al 2007, Gebhardt et al 2009, Jain et al 2011, Mylonakis et al 2011, Langhorn et al 2014). Investigations in people have shown that CRP might be a reliable marker for confirming the presence of early sepsis (Pancer et al 2011). However, CRP and SAA are not specific for detection of bacterial infection, because it is known to increase a variety of diseases (Mastrorilli et al 2007, Gebhardt et al 2009, Jain et al 2011).

In this study, PC activity significantly increased in dogs with PVE compared to healthy dogs (Table 1). Increased activity of PC could be related to insufficient development of hypercoagulable state that occurring disseminates intravascular coagulation or the disease started before 48 hours. Similarly, de Laforcade et al (2008) detected a significant decrease in PC activity from days 1 to 2, followed by a gradual rising in PC activity over time, and this PC activity may remain a useful marker of disease severity in dogs with sepsis. It was reported that PC activity highly decreased in dogs with sepsis (Yan and Dhainaut 2001, de Laforcade et al 2003) and dogs with systemic inflammatory syndrome (Bauer and Moritz 2013). However, Bauer and Moritz (2013) reported increases in PC activity in few dogs with systemic inflammatory response syndrome be interpreted as counter reaction following the activation of the coagulation system due to the inflammatory process. This result was not coincident the results of previous study (Yan and Dhainaut 2001, de Laforcade et al 2003), but this result coincident the results to reported de Laforcade et al (2008).





Cytokines play considerable role in the development and regulation of immune response, thus cytokine profiles contribute to the effect immunity level in the diseases (Ceremi 1992). IL-1 $\beta$ , TNF- $\alpha$  and INF- $\gamma$  are produced by inflammatory cells (Jain et al 2011). The release of proinflammatory cytokines such as IL-1 $\beta$ , IL-6 and TNF- $\alpha$  by monocytes/macrophage and activated T-lymphocyte is considered to be a key event in the development of sepsis (de Boer et al 1992). Bacterial toxins lead to inflammation mediators release from mononuclear phagocytes in sepsis. IL-1 $\beta$  and TNF- $\alpha$  are important inflammatory cytokines implicated in a variety of disease in dogs (Nakagawa-Tosa et al 1995, Alsemgeest et al 1996, Ohtsuka et al 1997, Baggio et al 2005). It was reported that IL-6 and TNF- $\alpha$  levels significantly elevated in sepsis, and these parameters might be useful markers for evaluating sepsis (Nemzek et al 2001, Rau et al 2007, Song et al 2012). Serum IL-1 $\beta$  and TNF- $\alpha$  concentrations usually increase in inflammatory disease of dogs (Miyomato et al 1996, Grone et al 2000). Fransson et al (2007) reported that IL-6, TNF $\alpha$  and CRP levels increased in dogs with pyometra and dogs with systemic inflammatory response syndrome. In this study, serum IL-1 $\beta$ , TNF- $\alpha$  and INF- $\gamma$  concentrations remarkably increased in dogs with PVE when compared to healthy dogs (Table 1). The reason of increase in IL-1 $\beta$  and TNF $\alpha$  levels could be related to release of inflammatory mediators against bacterial toxins which occurs in sepsis. Bacterial translocation and coliform septicemia are known to be common reasons in the mortality of dogs with PVE (Isogai et al 1989). Furthermore, INF- $\gamma$  is produced against viral infection (Niewold et al 2003). Therefore, selected cytokines such as IL-1 $\beta$ , TNF- $\alpha$  and INF- $\gamma$  might be useful indicators in assessing the clinical severity of sepsis in dogs. These results were coincident the results to reported of Miyomato et al (1996), Grone et al (2000), Nemzek et al (2007) and Rau et al (2007).

## Conclusion

Increased CRP, SAA, IL-1 $\beta$ , TNF- $\alpha$  and INF- $\gamma$  concentrations in dogs with PVE may be accepted to be significant markers of systemic inflammation in parvoviral infection in dogs.

## References

- Alsemgeest SP, van Klooster GA, van Miert AS, Hulskamp-Koch CK, Gruys E, 1996. Primary bovine hepatocytes in the study of cytokine induced acute-phase protein secretion in vitro. *Vet Immunol Immunopathol*, 53, 179-184.
- Baggio V, Ott F, Fischer RW, Gram H, Peele J, Spreng D, Sckmotel H, Jungi TW, 2005. Production of antibodies to canine IL-1 $\beta$  and canine TNF to assess the role of proinflammatory cytokines. *Vet Immunol Immunopathol*, 107, 27-39.
- Bauer N, Moritz A, 2013. Coagulation response in dogs with and without systemic inflammatory response syndrome - Preliminary results. *Res Vet Sci*, 94, 123-121.
- Bochsler PN, Slauson DO, 2002. Inflammation and repair tissue, in: *Mechanism of Disease. A Textbook of Comparative General Pathology*, third edition, Mosby, St. Louis, USA, pp: 140-245.
- Carry C, 2011. Acute phase proteins in animals. *Prog Mol Biol Transl Sci*, 105,113-150.
- Cary C, 2013. Biomarkers of inflammation in exotic pets. *J Exot Pet Med*, 22, 245-250.
- Ceremi A. 1992. Inflammatory cytokines. *Clin Immunol Immunopathol*, 62, 3-10.
- Ceron JJ, Eckersall PD, Martinez-Subiela S, 2005. Acute phase proteins in dogs and cats, current knowledge and future perspectives. *Vet Clin Pathol*, 34, 85-99.
- de Boer JP, Wolbink GJ, Thijs LG, Baars JW, Wagstaff J, Hack CE, 1992. Interplay of complement and cytokines in the pathogenesis of septic shock. *Immunopharmacology*, 24, 135-148.
- de Laforcade AM, Freeman LMS, Shaw SP, Brooks MB, Rozanski EA, Rush JE, 2003. Hemostatic changes in dogs with naturally occurring sepsis. *Vet Intern Med*, 17,674-679.
- de Laforcade AM, Rozanski EA, Freeman LM, Li W, 2008. Serial evaluation of protein C and antithrombin in dogs with sepsis. *J Vet Intern Med*, 22, 26-30.
- Eckersall PD, Bell R, 2010. Acute phase proteins, biomarkers of infection and inflammation in veterinary medicine. *Vet J*, 185, 23-27.
- Er C, Ok M, 2015. Levels of cardiac biomarkers and coagulation profiles in dogs with parvoviral enteritis. *Kafkas Univ Vet Fak Derg*, 21, 383-388.
- Fransson BA, Lagerstedt AS, Bergstrom A, Hagman R, Park JS, Evans MA, Ragle CA, 2007. C-reactive protein, tumor necrosis factor  $\alpha$  and interleukin-6 in dogs with pyometra and SIRS. *J Vet Emer Crit Care*, 17, 373-381.
- Gebhardt C, Hirschberger J, Rau S, Arndt G, Krainer K, Schweigert FJ, Brunnberg L, Kaspers B, Kohn B, 2009. Use of C-reactive protein to predict outcome in dogs with systemic inflammatory response syndrome or sepsis. *J Vet Emer Crit Care*, 19, 450-458.
- Goddard A, Leisewitz AL, 2010. Canine parvovirus. *Vet Clin Small Anim*, 40, 1041-1053.
- Grone A, Aldinger S, Baumgartner W, 2000. Interleukin-1 $\beta$ , -6, -12 and tumor necrosis factor- $\alpha$  expression in brain of dogs with canine distemper virus infection. *J Neuroimmunol*, 110, 20-30.
- Isogai E, Isogai H, Onuma M, Mizukoshi N, Hayashi M, Namioka S, 1989. Escherichia coli associated endotoxemia in dogs with parvovirus infection. *Nippon Jugaku Zasshi*, 51, 597-606.
- Jain S, Gautam V, Naseem S, 2011. Acute-phase proteins, as diagnostic tool. *J Pram BioallSci*, 1, 118-127.
- Jitpean S, Holst BS, Høglund OV, Pettersson A, Olsson U, Strage E, odersten F, Hagman R, 2014. Serum insulin-like factor-I, iron, C-reactive protein, and serum amyloid A for prediction of outcome in dogs with pyometra. *Theriogenology*, 30, 1-6.
- Kariuki NM, Nyaga PN, Buoro IBJ, Gathumbi PK, 1990. Effec-







- tiveness of fluids and antibiotics as supportive therapy of canine parvovirus 2 enterit in puppies. *Bull Anim Health Prod Afr*, 38, 379-389.
- Kjelgaard-Hansen M, Jacobsen S, 2011. Assay validation and diagnostic applications of major acute-phase proteins testing in companion animals. *Clin Lab Med*, 31, 51-70.
- Kocaturk M, Martinez S, Eralp O, Tvarijonaviciute A, Cero J, Yilmaz Z, 2010. Prognostic value of serum acute-phase proteins in dogs with parvoviral enteritis. *J Small Anim Pract*, 51, 478-483.
- Langhorn CMB, Goddard R, Andreasen A, Moldal EB, Tvarijonaviciute A, Kirpensteijn J, Jakobsen S, Persson F, Kjelgaard-Hansen M, 2014. Comparison of serum amyloid A and C-reactive protein as diagnostic markers of systemic inflammation in dogs. *Can Vet J*, 55, 161-168.
- Mastrorilli C, Dondi F, Agnoli C, 2007. Clinicopathologic features and outcome predictors of *Leptospira interrogans Australis* serogroup infection in dogs: A retrospective study of 20 cases (2001-2004). *J Vet Intern Med* 21, 3-10.
- McCaw DL, Hoskins JD, 2006. Canine viral enteritis, in: *Infectious Diseases of the Dogs and Cats*, Ed: Green CE, third edition, Saunders Elsevier, St. Louis, USA, pp: 63-73.
- Miyamota T, Fujinaga T, Yamashita K, Hagio M, 1996. Changes of serum cytokine activities and other parameters in dogs with experimentally induced endotoxic shock. *Jpn J Vet Res*, 44, 107-118.
- Murata H, Shimada N, Yoshioka M, 2004. Current research on acute phase proteins in veterinary diagnosis, an overview. *Vet J*, 168, 28-40.
- Mylonakis ME, Ceron JJ, Leontides L, Siarkou VI, Martinez S, Tvarijonaviciute A, Koutinas AF, Harrus S, 2011. Serum acute phase proteins as clinical phase indicators and outcome predictors in naturally occurring canine monocytic ehrlichiosis. *J Vet Intern Med*, 25, 811-817.
- Nakagawa-Tosa N, Morimatsu M, Kawasaki M, Nakatsuji H, Syuto B, Saito M, 1995. Stimulation of haptoglobin synthesis by interleukin-6 and tumor necrosis factor, but not by interleukin-1, in bovine primary cultured hepatocytes. *J Vet Med Sci*, 57, 219-223.
- Nemzek JA, Siddiqui J, Remick DG, 2001. Development and optimization of cytokine ELISAs using commercial antibody pairs. *J Immunol Methods*, 255, 149-157.
- Niewold TA, Tousaint MJM, Gruys E, 2003. Monitoring healthy by acute phase proteins Fourth European Colloquium on acute phase proteins. Segova, İspanya, pp: 57-67.
- Ohtsuka H, Ohki K, Tanaka T, Tajima M, Yoshino T, Takahashi K, 1997. Circling tumor necrosis factor and interleukin -1 after administration of LPS in adult cows. *J Vet Med Sci*, 19, 927-229.
- Ok M, Sen I, Birdane FM, Bektas HG, Turgut K, 2000. Diagnostic importance of ELISA and haemagglutination inhibition tests in canine parvoviral infection of dogs. *Indian Vet J*, 77, 465-467.
- Otto CM, Drobatz KJ, Soter C, 1997. Endotoxemia and tumor necrosis factor activity in dogs with naturally occurring parvoviral enteritis. *J Vet Intern Med*, 11, 65-70.
- Pancer G, Engelman E, Hoque F, Alam M, Rucinski J, Bernstein LH, 2011. C-reactive protein for the enhanced evaluation of the systemic inflammatory response syndrome (SIRS). *Open Clin Chem J*, 4, 1-9.
- Pollock RVH, Coyne MJ, 1993. Canine parvovirus. *Vet Clin North Am Small Anim Pract*, 23, 555-568
- Prittie J, 2004. Canine parvoviral enterit: A review of diagnosis, management and prevention. *J Vet Emerg Crit Care*, 14, 167-176.
- Rau S, Kohn B, Richter C, Fenske N, Kuchenhoff H, Hartmann K, Hartle S, Kaspers B, Hirschberger J, 2007. Plasma interleukin-6 response is predictive for severity and mortality in canine systemic inflammatory response syndrome and sepsis. *Vet Clin Pathol* 36, 253-260.
- Song R, Kim J, Yu D, Prak C, Park J, 2012. Kinetics of IL-6 and TNF $\alpha$  changes in canine models of sepsis induced by endotoxin. *Vet Immunol Immunopathol*, 146, 143-149.
- Turgut K, 2000. Disproteinemiler ve testler, in: *Veteriner Klinik Laboratuvar Teşhis*, ikinci baskı, Bahçivanlar Basım Sanayi, Konya, Türkiye, pp: 489-505.
- Yan SB, Dhainaut JF, 2001. Activated protein C versus protein C in severe sepsis. *Crit Care Med*, 29, 69-74.
- Yeşilbağ K, Yılmaz Z, Özkul A, Pratelli A, 2007. Etiological role of viruses in puppies with diarrhoea. *Vet Rec*, 161, 169-170.
- Yılmaz Z, Şentürk S, 2007. Characterization of lipid profiles in dogs with parvoviral enterit. *J Small Anim Pract*, 48, 643-650.
- Yule TD, Roth MB, Dreier K, Johnson AF, Palmer-Densmore M, Simmons K, Fanton R, 1997. Canine parvovirus vaccine elicits protection from the inflammatory and clinical consequences of the disease. *Vaccine*, 15, 720-729.